

Peer Reviewed

Title: Silicon uptake and translocation in plants

Author: Ma, Jian Feng, Okayama University

Publication Date: 04-14-2009

Publication Info:

The Proceedings of the International Plant Nutrition Colloquium XVI, Department of Plant Sciences, UC Davis, UC Davis

Permalink:

http://escholarship.org/uc/item/3pq8p5p0

Keywords:

Silicon, transporters, uptake, translocation, distribution, rice

Abstract:

Silicon (Si) is the second most abundant element after oxygen in the earth's crust and shows beneficial effects on plant growth and production by alleviating both biotic and abiotic stresses including diseases, pests, lodging, drought, and nutrient imbalance. Transporters involved in uptake and translocation of Si have been identified in Si-accumulating species including rice, barley and maize. Two kinds of transporters (influx and efflux) are involved in the Si uptake by the roots. The influx transporters (low silicon 1, Isi1) belongs to a Nod26-like major intrinsic protein (NIP) subfamily in aquaporin, while the efflux transporter (Lsi2) belongs to putative anion transporters. They are mainly expressed in the roots. Knockout of either Lsi1 or Lsi2 in rice results in defect of Si uptake, indicating that coupling of Lsi1 with Lsi2 is required for efficient uptake of Si in rice. Although the function of Lsi1 and Lsi2 is similar in rice, barley and maize, they show different expression pattern and cell-type specificity of localization. These differences may explain the differential Si uptake capacity among these species. On the other hand, Lsi6 is localized at the adaxial side of the xylem parenchyma cells in the leaf sheath and leaf blades of rice and maize. Knockout of rice Lsi6 does not affect the uptake of Si by the roots, but results in disordered deposition of silica in the shoots and increased excretion of Si in the guttation fluid, indicating that Lsi6 is a transporter responsible for the unloading of Si from the xylem.



eScholarship provides open access, scholarly publishing services to the University of California and delivers a dynamic research platform to scholars worldwide.

Introduction

Silicon (Si) is the second most abundant element after oxygen in the earth's crust and all plants contain Si in their tissues. Since there is still no evidence showing that Si is involved in the metabolism of plants, Si has not been recognized as an essential element according to the criteria of essentiality established by Arnon and Stout (1939). However, beneficial effects of Si have been observed in a wide variety of plant species. The characteristic function of Si is to help plants to overcome multiple biotic and abiotic stresses (Ma, 2004; Ma and Yamaji, 2006; Ma and Yamaji, 2008). For example, Si increases the resistance of plants to pests and pathogens. Silicon also enhances the resistance to drought and heavy-metal toxicity. Based on such important roles of Si in plants, Epstein and Bloom (2003) have proposed Si as "quasi-essential" for the growth of higher plants. In Japan, Si has been recognized as an "agronomically essential element" for rice, a typical Si-accumulating plant, because high accumulation of Si is required for optimal growth and sustainable production of rice.

Silicon is taken up by the roots in the form of silicic acid, a non-charged molecule. It is the same form to be translocated in the xylem from the roots to the shoots. In the shoot, with water loss through transpiration, silicic acid is concentrated and polymerized to silica (SiO_2) and finally deposited on the different tissues. Therefore, for the transfer of Si from the external solution (soil solution) to the different tissues, different transporters are required. Recently, great progresses have been made in identification of Si transporters involved in the uptake, translocation and distribution of Si.

Transporters Involved in Si Uptake

Radial transport of Si from the external solution into the root cells includes influx and efflux transporters.

Influx Si transporters

The first Si transporter (Lsi1) was identified from rice (Ma et al., 2006). Rice is able to accumulate Si over 10% in the shoots. By using a mutant approach, the gene *Lsi1* (Low silicon 1) was isolated by map-based cloning method. Lsi1 belongs to a Nod26-like major intrinsic protein (NIP) subfamily of aquaporin-like proteins. The predicted amino acid sequence has six transmembrane domains and two Asn-Pro-Ala (NPA) motifs, which are well conserved in typical aquaporins. By using a *Xenopus* oocyte assay system, Lsi1 shows an influx transport activity for silicic acid (Ma et al., 2006). The transport activity is not affected by low temperature (Mitani et al., 2008). Recently, it was reported that Lsi1 is also

permeable to arsenite (Ma et al., 2008).

The gene *Lsil* is constitutively expressed in the roots, but its expression is decreased to one fourth by Si supply (Ma et al., 2006). Within a root, the expression of *Lsil* was much lower in the root tip region between 0-10 mm than that in the basal regions (>10 mm) (Yamaji and Ma, 2007). Silicon uptake in the root tip region (0 to 10 mm) comprising both the apical meristem and the elongation zone is also much lower than that in the basal regions (>10 mm from the root tips). These observations indicate that the site of Si uptake is located in the mature regions of the roots rather than the root tips.

Investigation of rice *Lsi1* expression at different growth stages showed that the expression was transiently enhanced around the heading stage (Yamaji and Ma, 2007). A previous study showed that 67% of total Si was taken up during the reproductive stage from panicle initiation to heading in rice (Ma et al., 1989). Deficiency of Si during this stage results in a significant reduction in the grain yield, suggesting that a high Si uptake during this period is required for producing a high yield. Therefore, the increased expression of *Lsi1* during the heading stage coincides with a high Si requirement during this growth stage.

Lsi1 is localized in the main and lateral roots, but not in root hairs (Ma et al., 2006). This is consistent with the results of a previous physiological study that root hairs do not play any demonstrable role in silicon uptake, but that lateral roots contribute significantly to silicon uptake (Ma et al., 2001). Furthermore, in rice roots including seminal, lateral and crown roots, the Lsi1 protein is localized at both exodermis and endodermis, where Casparian strips exist. Interestingly, Lsi1 shows polar localization at the distal side of both the exodermis and endodermis cells (Fig. 1). Taken together, Lsi1 is responsible for the transport of silicic acid from the external solution to the root cells in rice.

Following identification of rice Lsi1, Si influx transporters have also been identified in maize (ZmLsi1) and barley (HvLsi1) (Chiba et al., 2009; Mitani et al., 2009). At amino acid level ZmLsi1 and HvLsi1 show 82% and 82% identity with OsLsi1, respectively. Both HvLsi1 and ZmLsi1 also show Si influx transport activity like OsLsi1, but their cell-type specificities of localization and expression patterns are different from OsLsi1. HvLsi1 and ZmLsi1 are localized at the epidermal, hypodermal and cortical cells (Chiba et al., 2009; Mitani et al., 2009). Furthermore, the expression levels of both *HvLsi1* and *ZmLsi1* are unaffected by Si. These differences may be related to the root structures and Si uptake ability among different species.

Efflux transporters of silicon

The efflux transporter gene (*Lsi2*) of Si was also cloned using a novel rice mutant (*lsi2*) defective in Si uptake (Ma et al., 2007). This gene *Lsi2* is predicted to encode a membrane protein with 11 transmembrane domains, belonging to a putative anion transporter without any similarity with the silicon influx transporter Lsi1. In *Xenopus laevis* oocytes, Lsi2 does

not show influx transport activity for silicic acid but shows efflux transport activity (Ma et al., 2007). These results suggest that in contrast to the influx transporter Lsi1, Lsi2 is a silicon efflux transporter, which is capable of transporting silicon out of the cells. Interestingly, the efflux of Si was inhibited by a low temperature treatment and by three protonophores; 2,4-dinitrophenol (DNP), carbonylcyanide 3-chlorophenylhydrazone (CCCP) and carbonylcyanide p-(trifluoromethoxy)penylhydrazone (FCCP). Furthermore, the efflux activity of Lsi2 is increased at lower external pH values (Ma et al., 2007). All these results suggest that transport of silicon by Lsi2 is an energy-dependent active process, which is driven by the proton gradient.

Lsi2 is mainly expressed in the roots as is *Lsi1*. The mRNA accumulation was constitutive but it decreased to one fourth by continuous silicon supply (Ma et al., 2007). Furthermore, there is little accumulation of *Lsi2* transcripts in the root tip (0-10 mm), but much accumulation in the mature parts of the roots. These expression patterns are similar to those of *Lsi1* (Ma et al., 2006, Yamaji and Ma, 2007), suggesting that the expression of *Lsi1* and *Lsi2* may be regulated in a similar manner. Comparison of the promoter region revealed common domains in the *Lsi1* and *Lsi2* (Ma et al., 2007).

Like Lsi1, Lsi2 is also localized at the exodermis and the endodermis cells of the roots. However, in contrast to Lsi1 localized on the distal side, Lsi2 was localized on the proximal side of the exodermis and the endodermis cells. Si efflux transporters in other plant species have not been identified.




Figure 1. Silicon transporters involved in the uptake and distribution of Si in rice.

An efficient transport system for Si in rice by coupling of Lsi1 and Lsi2

As described above, both Si transporters (Lsi1 and Lsi2) are localized on the exodermis and endodermis, where Casparian strips exist (Fig. 1). Casparian strips prevent solutes to pass freely from the external solution to the stele. In fact, silicon is highly deposited on the exodermis and endodermis. Both Lsi1 and Lsi2 are required for transcellular transport of silicon to the stele. Moreover, rice roots are characterized by the formation of aerenchyma, which is accompanied by the destruction of almost all cortex cells except the exodermis and the endodermis. Therefore, Si transported into the exodermis cells by Lsi1 is released by Lsi2 into the apoplast of a spoke-like structure across the aerenchyma. Silicon is then transported into the endodermis cells by Lsi2 (Fig.1). Coupling of Lsi1 and Lsi2 in the same cell in the Casparian strips is required for efficient transport of Si across the cells into the stele.

A recent study showed that genotypic differences in the Si accumulation in rice can result from the difference in the expression of Si transporter genes in roots (Ma et al., 2007). The expression of both Lsi1 and Lsi2 was lower in a variety with low Si uptake capacity than in a variety with high Si uptake capacity. However, there is no difference in the cellular localization of these two transporters between low and high Si uptake varieties. Therefore, high expression of both Lsi1 and Lsi2 is required to enhance Si uptake.

Transporters Involved in Translocation and Distribution of Si

Silicon transported by Lsi1 and Lsi2 into the stele is then translocated to the shoot by transpirational flow through the xylem. More than 90% of Si taken up by the roots is translocated to the shoots (Ma and Takahashi, 2002). Silicon is present in the xylem at high concentrations and in the form of monosilicic acid (Casey et al., 2003; Mitani et al., 2005). Recently, a transporter (Lsi6), which is responsible for the export of silicic acid from the xylem and for the subsequent distribution of Si, was identified in rice.

Lsi6 is a homolog of Lsi1. It also shows transport activity for silicic acid in the oocyte assay. However, different from Lsi1 and Lsi2, Lsi6 is also expressed in the leaf sheath and leaf blades in addition to the root tips (Yamaji et al., 2008). Lsi6 is localized in the adaxial side of the xylem parenchyma cells in the leaf sheath and leaf blades (Fig. 1). Knockout of Lsi6 does not affect the uptake of Si by the roots, but affects the silica deposition pattern in the leaf blades and sheaths. The density of silicified dumbbell-shape and motor cells in the knockout line is decreased compared with wild-type rice. The abaxial epidermis cells is observed to be silicified in the mutant, but infrequently in the wild-type rice (Yamaji et al., 2008). Furthermore, knockout of Lsi6 results in an increased excretion of Si in the guttation fluid. These results suggest that knockout of *Lsi6* resulted in alteration of the Si pathway to

the specific cells. Therefore, cell-type specific silicification depends on the symplastic pathway of Si delivered by Lsi6 rather than apoplastic pathway. A similar transporter ZmLsi6 has also been identified in maize (Mitani et al., 2009).

The level of Si accumulation in the shoots greatly differ among plant species, ranging from 0.1% to 10% of dry weight (Hodson et al., 2005; Ma and Takahashi, 2002). However, the molecular mechanisms for these differences are still unknown. Identification of more Si transporters in different plant species in future, will help to better understand the mechanisms of Si uptake, translocation and distribution in plants.

ACKNOWLEDGMENTS

This work was supported by the Program of Promotion of Basic Research Activities for Innovative Biosciences (BRAIN), a Grant-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Culture, Sports, Science and Technology of Japan (No. 17078008) and a grant from the Ministry of Agriculture, Forestry and Fisheries of Japan (Genomics for Agricultural Innovation IPG-006) to J. F. M.

References

Arnon, D.I. and Stout, P.R. (1939) The essentiality of certain elements in minute quantity for plants with special reference to copper. Plant Physiol., **14**: 371-375

Casey, W.H., Kinrade, S.D., Knight, C.T.G, Rains, D.W., and Epstein, E. (2003) Aqueous silicate complexes in wheat, *Triticum aestivum* L. Plant Cell Environ. **27**: 51-54.

Chiba, Y., Mitani, N., Yamaji, N., and Ma, J.F. (2009) HvLsi1 is a silicon influx transporter in barley. Plant J. 57: 810-818.

Epstein, E. and Bloom, A. J. (2003) Mineral Nutrition of Plants: Principles and Perspectives, Second Edition, John Wiley & Sons, New York.

Hodson, MJ., White, PJ., Mead, A., and Broadley, MR. (2005) Phylogenetic variation in the silicon composition of plants. Ann. Bot. **96:** 1027-1046

Ma, J. F., Nishimura, K. and Takahashi, E. (1989) Effect of silicon on the growth of rice plant at different growth stages. Soil Sci. Plant Nutri. **35**: 347-356

Ma, J. F., Goto, S., Tamai, K and Ichii, M. (2001) Role of root hairs and lateral roots in silicon uptake by rice. Plant Physiol. 127: 1773-1780

Ma, J. F. and Takahashi, E. (2002) Soil, Fertilizer, and Plant Silicon Research in Japan. first edition, Elsevier Science, Amsterdam, 2002.

Ma, J.F. (2004). Role of silicon in enhancing the resistance of plants to biotic and abiotic stresses. Soil Sci. Plant Nutr. **50**: 11-18.

Ma, J.F., Tamai, K., Yamaji, N., Mitani, N., Konishi, S., Katsuhara, M., Ishiguro, M., Murata, Y., and Yano, M. (2006) A silicon transporter in rice. Nature **440**: 688-691.

Ma, J.F., and Yamaji, N. (2006) Silicon uptake and accumulation in higher plants. Trends

Plant Sci. 11: 392-397.

Ma, J.F., and Yamaji, N. (2008) Functions and transport of silicon in plants. Cell Mol. Life Sci. 65: 3049-3057.

Ma, J.F., Yamaji, N., Mitani, N., Tamai, K., Konishi, S., Fujiwara, T., Katsuhara, M., and Yano, M. (2007). An efflux transporter of silicon in rice. Nature **448**: 209-212.

Ma, J. F., Yamaji, N., Tamai, K., and Mitani, N. (2007) Genotypic difference in Si uptake and expression of Si transporter genes in rice. Plant Physiol. **145** : 919-924.

Ma, J. F., Yamaji, N., Tamai, K. and Mitani, N. (2007) Genotypic difference in Si uptake and expression of Si transporter genes in rice. Plant Physiol. **145** : 919-924

Ma, J. F., Yamaji, N., Mitani, N., Xu, X. Y., Su, Y. H., McGrath, S. and Zhao, F. J. (2008) Transporters of arsenite in rice and their role in arsenic accumulation in rice grain. Proc. Nat. Acad. Sci. 105: 9931-9935

Mitani, N., Ma, J. F., and Iwashita, T. (2005) Identification of the silicon form in xylem sap of rice (*Oryza sativa* L.). Plant Cell Physiol. **46:** 279-283.

Mitani, N., Yamaji, N. and Ma, J. F. (2008) Characterization of substrate specificity of a rice silicon transporter, Lsi1. *Pflugers Arch*-Eur. J. Physiol. **456**: 679-686.

Mitani, N., Yamaji, N., and Ma, J.F. (2009) Identification of maize silicon influx transporters. Plant Cell Physiol. 50: 5-12.

Yamaji, N., and Ma, J.F. (2007) Spatial distribution and temporal variation of the rice silicon transporter Lsi1. Plant Physiol. **143**: 1306-1313.

Yamaji, N., Mitatni, N., and Ma, J.F. (2008). A transporter regulating silicon distribution in rice shoots. Plant Cell, **20**: 1381-1389.