

This article was downloaded by: [Universidad Publica de Navarra]

On: 16 May 2011

Access details: Access Details: [subscription number 936140347]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Plant Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597277>

### NITRATE/AMMONIUM RATIOS AFFECT RYEGRASS GROWTH AND NITROGEN ACCUMULATION IN A HYDROPONIC SYSTEM

Haiqin Cao<sup>a</sup>; Ying Ge<sup>a</sup>; Dong Liu<sup>a</sup>; Qianjin Cao<sup>a</sup>; Scott X. Chang<sup>b</sup>; Jie Chang<sup>a</sup>; Xiulin Song<sup>a</sup>; Xue Lin<sup>a</sup>

<sup>a</sup> College of Life Sciences, Zhejiang University, Hangzhou, China <sup>b</sup> Department of Renewable Resources, University of Alberta, Edmonton, Alberta, Canada

Online publication date: 01 December 2010

**To cite this Article** Cao, Haiqin , Ge, Ying , Liu, Dong , Cao, Qianjin , Chang, Scott X. , Chang, Jie , Song, Xiulin and Lin, Xue(2011) 'NITRATE/AMMONIUM RATIOS AFFECT RYEGRASS GROWTH AND NITROGEN ACCUMULATION IN A HYDROPONIC SYSTEM', Journal of Plant Nutrition, 34: 2, 206 – 216

**To link to this Article:** DOI: 10.1080/01904167.2011.533322

**URL:** <http://dx.doi.org/10.1080/01904167.2011.533322>

## PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## NITRATE/AMMONIUM RATIOS AFFECT RYEGRASS GROWTH AND NITROGEN ACCUMULATION IN A HYDROPONIC SYSTEM

Haiqin Cao,<sup>1</sup> Ying Ge,<sup>1</sup> Dong Liu,<sup>1</sup> Qianjin Cao,<sup>1</sup> Scott X. Chang,<sup>2</sup> Jie Chang,<sup>1</sup> Xiulin Song,<sup>1</sup> and Xue Lin<sup>1</sup>

<sup>1</sup>College of Life Sciences, Zhejiang University, Hangzhou, China

<sup>2</sup>Department of Renewable Resources, University of Alberta, Edmonton, Alberta, Canada

□ *Ryegrass has increasingly been used in constructed wetlands for treatment of eutrophic wastewater. To properly match plant species with the characteristics of wastewater being treated, it is important to know the performance of plant species under different nitrate/ammonium ( $\text{NO}_3^-/\text{NH}_4^+$ ) ratios. We investigated ryegrass (*Lolium perenne* L.) dry matter (DM) production and N accumulation under five  $\text{NO}_3^-/\text{NH}_4^+$  ratios (100/0, 75/25, 50/50, 25/75, 0/100) in a hydroponic system. The results showed that ryegrass total DM, shoot DM, root DM and nitrogen (N) accumulation were greater under  $\text{NO}_3^-/\text{NH}_4^+$  ratios of 50/50 and 75/25 than under other  $\text{NO}_3^-/\text{NH}_4^+$  ratios, indicating that ryegrass can be best used in constructed wetlands for treating wastewater with such  $\text{NO}_3^-/\text{NH}_4^+$  ratios to achieve high biomass production and efficient removal of N. On the other hand, for treating wastewater with either  $\text{NO}_3^-$  or  $\text{NH}_4^+$  dominate the inorganic N, other plant species that are more adapted to such conditions should be explored.*

**Keywords:** wastewater, N concentration, dry matter production, root growth, root shoot ratio

### INTRODUCTION

Excessive discharge of nutrients into rivers may cause eutrophication and deteriorate ecosystem health (Tylova et al., 2005). In many small cities in China, sewage is still being directly drained into water bodies without treatment (Liu et al., 2009), which has caused serious water pollution and intensified water shortage. Therefore, proper treatment of eutrophic wastewater is crucial for maintaining a healthy environment, clean water supply, and human health. Constructed wetlands (CWs) have emerged as a viable wastewater treatment technology (Liu et al., 2009). In CWs, plant

Received 21 January 2009; accepted 29 March 2009.

Address correspondence to Ying Ge, College of Life Sciences, Zhejiang University, 388 Yuhangtang Road, Hangzhou 310058, P. R. China. E-mail: geying@zju.edu.cn

nitrogen (N) uptake and denitrification by microorganisms are the two main mechanisms responsible for N removal. Between the two mechanisms, plant uptake of N usually plays a minor role, accounting for 5–23% of the total N removal (Ge et al., 2007). However, plants can greatly contribute to N removal by affecting nitrification and denitrification intensity in the root zone (Stottmeister et al., 2003; Ge et al., 2007). For instance, plant roots can release organic compounds and provide a suitable environment for the microorganisms (Sundaravadivel and Vigneswaran, 2001). In order to achieve efficient N removal from wastewater, plants with high biomass production, especially root biomass production, should be given priority.

Wastewater from different sources has different characteristics, such as different nitrate ( $\text{NO}_3^-$ )/ammonium ( $\text{NH}_4^+$ ) ratios. The inorganic N composition in wastewater typically has between 0.4 and 99.1%  $\text{NH}_4^+$  and between 0.9 and 99.6%  $\text{NO}_3^-$  (Liu et al., 2009). Plant species may have a preference between  $\text{NO}_3^-$  and  $\text{NH}_4^+$ , so species should be carefully selected according to the characteristics of the wastewater (Sundaravadivel and Vigneswaran, 2001). Different  $\text{NO}_3^-/\text{NH}_4^+$  ratios in the N supply can affect the rate of plant growth as well as biomass allocation between shoot and root (Guo et al., 2002; Bruck and Guo, 2006). Most species grow better and accumulate more N when supplied with a mixture of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  (Ali et al., 2001; Guo et al., 2002). Species such as tea (*Camellia sinensis*) prefer  $\text{NH}_4^+$  as the N-source (Ruan et al., 2007), whereas wheat (*Triticum* spp.) is tolerant to  $\text{NH}_4^+$  but only to low  $\text{NH}_4^+$  concentrations (Cox and Reisenauer, 1973). Many other plants are highly sensitive to  $\text{NH}_4^+$  (Ali et al., 2001; Britto and Kronzucker, 2002). For such species, when  $\text{NH}_4^+$  is supplied as the sole N source, plants can develop symptoms of toxicity and the root growth can be severely impaired (Lasa et al., 2001; Britto and Kronzucker, 2002).

However, most relevant studies have been focused on cultivated plants (Ali et al., 2001; Lasa et al., 2001; Bruck and Guo, 2006; Ruan et al., 2007), especially vegetables, with emphasis on the economic yield and inorganic N concentrations in plant tissues (which has implications for human health). Although the preference of *Phragmites* and *Glyceria*, two species commonly used in CWs, for  $\text{NH}_4^+$  vs  $\text{NO}_3^-$  have been studied by Tylova et al. (2005), the preference for  $\text{NH}_4^+$  vs  $\text{NO}_3^-$  among other plant species used in CWs for wastewater treatment is poorly understood.

Ryegrass (*Lolium perenne* L.) is a species commonly used in vertical flow CWs (Chang et al., 2004). It can accumulate considerable amounts of nutrients for removal through harvesting (Caicedo et al., 2000). But no detailed studies on the effects of different inorganic N forms on ryegrass for wastewater treatment have been found in the literature. In this study, a hydroponic system was used in order to minimize the transformation of N between the two inorganic forms by microorganisms. The objectives of this study were to (1) determine the impact of  $\text{NO}_3^-/\text{NH}_4^+$  ratio on total and root biomass production of ryegrass; and (2) investigate the effects of  $\text{NO}_3^-/\text{NH}_4^+$  ratio

on nitrogen uptake and thus nitrogen removal from the culture solution by ryegrass. Results from this study can provide information for ryegrass deployment in CWs.

## MATERIALS AND METHODS

### Plant Materials and Growth Conditions

The experiment was conducted in the greenhouse (120°05' E, 30°18' N) in the College of Life Sciences, Zhejiang University, Hangzhou, in south-east China. Uniform ryegrass seeds were selected and sown on a tray to germinate in distilled water. After about ten days when the first true leaf emerged, ryegrass seedlings were selected for uniformity and transplanted to plastic pots filled with 2 L nutrient solutions with different inorganic N compositions ( $\text{NO}_3^-/\text{NH}_4^+$  ratios of 100/0, 75/25, 50/50, 25/75, 0/100) at a constant N concentration of 8 mmol L<sup>-1</sup>. All treatments were replicated four times, for a total of 20 pots in the experiment, with each pot containing six plants. The basal nutrient solution was a modified Hoagland's nutrient solution (Hoagland and Arnon, 1950) that contained the following macronutrients (in mmol L<sup>-1</sup>): potassium ( $\text{K}^+$ ) 7, magnesium ( $\text{Mg}^{2+}$ ) 2, phosphate ( $\text{PO}_4^{3-}$ ) 1, and calcium ( $\text{Ca}^{2+}$ ) 5 (Table 1). Micronutrients were supplied as follows: 2.86 mg of boric acid ( $\text{H}_3\text{BO}_3$ ), 0.08 mg of copper sulfate ( $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ ), 0.22 mg of zinc sulfate ( $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ ), 1.81 mg of manganese chloride ( $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ ), 0.09 mg of molybdic acid ( $\text{H}_2\text{MoO}_4 \cdot 4\text{H}_2\text{O}$ ), and 7.645 mg iron (Fe)- ethylenediaminetetraacetic acid (EDTA) per liter of water (Zhang et al., 2005). The pH of the solution was adjusted to  $6.0 \pm 0.2$  with dilute sodium hydroxide (NaOH) or hydrochloric acid (HCl). Plants were grown with a 14 h photoperiod, at day/night temperatures of 25/15°C and 70–80% relative humidity at an irradiance of *ca.* 280  $\mu\text{mol photon}$

**TABLE 1** Compositions (mmol L<sup>-1</sup>) of nutrient solutions with different  $\text{NH}_4^+/\text{NO}_3^-$  ratios but the same N concentration (8 mmol L<sup>-1</sup>)

Nutrient source	$\text{NO}_3^-/\text{NH}_4^+$ ratio				
	100/0	75/25	50/50	25/75	0/100
	mmol L <sup>-1</sup>				
$\text{KNO}_3$	4.8	2.8	0.8	2.0	0.0
$\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$	1.6	1.6	1.6	0.0	0.0
$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	3.4	3.4	3.4	5.0	5.0
$(\text{NH}_4)_2\text{SO}_4$	0.0	1.0	2.0	3.0	4.0
$\text{KH}_2\text{PO}_4$	1.0	1.0	1.0	1.0	1.0
$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	2.0	2.0	2.0	2.0	2.0
KCl	1.2	3.2	5.2	4.0	6.0

$\text{m}^{-2} \text{s}^{-1}$ . The volume of culture solution in each pot was maintained at the initial volume of 2 L by adding distilled water every two days. Nutrient solutions were replaced every 10 days.

### Measurement and Calculation

After twenty days of growth, fully expanded leaves were selected to determine leaf chlorophyll concentrations. Chlorophyll was extracted using an ethane and ethanol mixture at a volume ratio of 2:1 according to Peng and Liu (1992). Chlorophyll concentration was determined spectrophotometrically in the clear supernatant by measuring the absorbance at 663 and 645 nm for chlorophyll a and b, respectively, using a HP 751 (Hewlett Packard, Shanghai, China) spectrophotometer. Chlorophyll concentrations were calculated using the equations given in Peng and Liu (1992). Before harvest, the rate of net photosynthesis was measured with a portable photosynthesis system (Licor-6400, Licor, Lincoln, NE, USA). Prior to the measurement, the leaves were illuminated at approximately  $600 \mu\text{mol photon m}^{-2} \text{s}^{-1}$  for 30 min in a growth chamber.

After 40 days of growth, ryegrass was harvested and divided into shoots (ryegrass has only leaves and root, no stem) and roots. The length of the longest root for each plant was measured. All leaves from each plant were collected and scanned with a scanner (ScanMaker 4900, Microtek International Inc., Cerritos, CA, USA) immediately after harvest. The WinFLORA Pro 2002a software (Regent Instruments Inc., Quebec, Canada) was used to determine leaf area (LA) from the images scanned. Plant shoot and root were dried in a forced air oven at  $65^\circ\text{C}$  for 72 h to determine dry matter (DM) yield, referred to as shoot DM, root DM and total DM (the sum of shoot and root biomass). The dried plant material was ground and digested to determine  $\text{NH}_4^+$  and  $\text{NO}_3^-$  concentrations in shoots and roots. From each sample, 0.2 g of material was weighed to analyze nutrient concentrations. The samples were digested by the Kjeldahl method applying sulfuric acid ( $\text{H}_2\text{SO}_4$ ) + hydrogen peroxide ( $\text{H}_2\text{O}_2$ ). The  $\text{NH}_4^+$  and  $\text{NO}_3^-$  concentrations were determined photometrically at wavelength of 420 nm and 210 nm, respectively (Lu et al., 2004). And total N concentration (TN) equals the sum of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  concentrations.

The following biometric characteristics were calculated following Hunt (1978): root:shoot ratio (R/S, calculated by dividing root DM by shoot DM), and specific leaf area (SLA, leaf area divided by leaf DM). Nitrogen uptake was calculated based on the biomass and concentration data.

### Statistical Analysis

Analysis of variance (ANOVA) was performed using the general linear model univariate procedure in SPSS (SPSS 15.0, SPSS Inc., Chicago, IL,

USA). The least significant difference was performed on each variable when there was a significant treatment effect from the ANOVA analysis, statistical significance was determined at  $\alpha = 0.05$ . All data are presented as mean  $\pm$  standard error (S.E.).

## RESULTS

### Biomass Production and Allocation

The  $\text{NH}_4^+$  to  $\text{NO}_3^-$  ratio affected the shoot and root growth of ryegrass. Total DM and shoot DM of ryegrass plants were higher under 75/25 and 50/50  $\text{NO}_3^-/\text{NH}_4^+$  ratios than under the other three treatments ( $P < 0.05$ ). In addition, total DM or shoot DM were not different between the 100%  $\text{NO}_3^-$  and 100%  $\text{NH}_4^+$  treatments ( $P > 0.05$ ) (Table 2). The ammonium only treatment resulted in root DM on average 68 and 61% lower than the 75/25 and 50/50 treatments ( $P < 0.05$ ), respectively (Table 2). We also found that root DM under the  $\text{NO}_3^-$  only treatment was also significantly lower than that under the 75/25 treatment, but obviously higher than that under the  $\text{NH}_4^+$  only treatment ( $P < 0.05$ ). A near linear decrease of R/S ratio was observed when the proportion of  $\text{NH}_4^+$  in the nutrient solution increased (Table 2). The effect of N form on the partition of assimilates between shoot and root is demonstrated in our experiment: only 4.5% of the total DM was allocated to the roots exposed to 100%  $\text{NH}_4^+$ .

### Physiological and Morphological Traits

No difference was found in net carbon dioxide ( $\text{CO}_2$ ) assimilation rate per unit leaf area among the five treatments (Table 2). Leaf area was higher under 75/25 and 50/50 ratios than under 100/0, 25/75 and 0/100 ratios ( $P < 0.05$ ). The chlorophyll concentration increased with the increase in the proportion of  $\text{NH}_4^+$  in solutions, while the reverse was true for SLA. Root length of plants decreased with the increasing proportion of  $\text{NH}_4^+$ . When  $\text{NH}_4^+$  reached 100%, root length was the lowest compared with plants in other treatment groups containing  $\text{NO}_3^-$  ( $P < 0.05$ ) (Table 2).

### Tissue N Concentration and Accumulation


A significant increase in root nitrate concentration in plants fed with  $\text{NO}_3^-$  alone over the other treatments was observed ( $P < 0.05$ , Figure 1). Generally, the greater the concentration of  $\text{NH}_4^+$  in the culture solution, the greater N concentration in plant shoots and roots. N concentration reached peak in plants grown with 100%  $\text{NH}_4^+$  (Figure 1).

Nitrogen accumulation mostly occurred in shoots, which accounted for over 90% of the total N accumulation (Figure 2). Ryegrass grown under


**TABLE 2** Shoot, root and total dry matter (DM), root/shoot ratio (R/S), leaf area (LA), chlorophyll concentration (Chl), net CO<sub>2</sub> assimilation rate ( $P_n$ ) of ryegrass (*Lolium perenne* L.) grown under five different NO<sub>3</sub><sup>-</sup>/NH<sub>4</sub><sup>+</sup> ratios. Values given are mean ± S.E. (n = 4)

Plant trait	NO <sub>3</sub> <sup>-</sup> /NH <sub>4</sub> <sup>+</sup> ratio				
	100/0	75/25	50/50	25/75	0/100
Shoot DM (g plant <sup>-1</sup> )	1.28 ± 0.08 b	2.22 ± 0.22 a	2.43 ± 0.11 a	1.42 ± 0.29 b	1.47 ± 0.18 b
Root DM (g plant <sup>-1</sup> )	0.14 ± 0.02 bc	0.22 ± 0.02 a	0.18 ± 0.02 ab	0.11 ± 0.01 cd	0.07 ± 0.01 d
Total DM (g plant <sup>-1</sup> )	1.5 ± 0.07 b	2.44 ± 0.24 a	2.61 ± 0.11 a	1.55 ± 0.31 b	1.57 ± 0.22 b
R/S	0.11 ± 0.02 a	0.1 ± 0.01 ab	0.07 ± 0.01 b	0.08 ± 0.01 b	0.04 ± 0.01 c
LA (cm <sup>2</sup> )	603.73 ± 47.04 b	972.2 ± 51.65 a	1032.16 ± 39.71 a	599.90 ± 98.11 b	546.03 ± 44.51 b
Chl (mg g <sup>-1</sup> )	1.59 ± 0.12 b	2.04 ± 0.1 ab	2.23 ± 0.08 a	2.26 ± 0.19 a	2.50 ± 0.27 a
$P_n$ (μmol CO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup> )	13.84 ± 0.87 a	13.08 ± 1 a	12.89 ± 0.81 a	14.54 ± 0.94 a	12.45 ± 0.54 a
SLA (cm <sup>2</sup> g <sup>-1</sup> )	471.46 ± 14.83 a	437.56 ± 17.66 a	424.54 ± 22.58 ab	378.23 ± 15.32 bc	347.68 ± 9.75 c
Root length (cm)	26.92 ± 2.03 a	27.5 ± 1.65 a	26.28 ± 1.39 ab	21.99 ± 1.96 bc	20.07 ± 0.5 c


Treatment means with different lowercase letters are significantly different ( $P < 0.05$ ) for each listed parameter.



**FIGURE 1** Concentrations of TN (open columns) and NO<sub>3</sub><sup>-</sup> (dashed columns) in shoots and roots of ryegrass (*Lolium perenne* L.) grown in solutions with different NO<sub>3</sub><sup>-</sup>/NH<sub>4</sub><sup>+</sup> ratios. Values given are mean ± S.E. (n = 4). Values with different letters are significantly different (P < 0.05) among the treatments for each parameter measured either for root or shoot.



**FIGURE 2** N accumulation in shoots (open columns) and roots (solid columns) of ryegrass grown in solutions with different NO<sub>3</sub><sup>-</sup>/NH<sub>4</sub><sup>+</sup> ratios. Values given are mean ± S.E. (n = 4). Values with different letters are significantly different (P < 0.05) among the treatments for either root or shoot.



**FIGURE 3** Relationships between plant dry mass and N accumulation in ryegrass. [(a) shoot, and (b) root] and relationships between N concentration and N accumulation in ryegrass [(c) shoot, (d) root].

$\text{NO}_3^-/\text{NH}_4^+$  ratios of 75/25 and 50/50 had the greatest shoot N accumulation, while the 100%  $\text{NH}_4^+$  had the intermediate and 100%  $\text{NO}_3^-$  had the lowest shoot N accumulation. Positive correlations were found between shoot DM and shoot N accumulation (Figure 3a) and between root DM and root N accumulation (Figure 3b). However, N concentration was not correlated with N accumulation in shoot or root (Figures 3c and 3d).

## DISCUSSION

### Effects of $\text{NO}_3^-/\text{NH}_4^+$ Ratios on Plant Biomass Production

Many previous studies have reported that a mixture of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  supply is better for plant growth compared with  $\text{NH}_4^+$  or  $\text{NO}_3^-$  alone (Ali et al., 2001; Guo et al., 2002; Zhang et al., 2007). In this study total and shoot DM yields were the highest in 75/25 and 50/50 treatments, suggesting that ryegrass prefers close to equal amounts of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  supply in the growth media, with slight preference for  $\text{NO}_3^-$ . Furthermore, the solutions with  $\text{NO}_3^-/\text{NH}_4^+$  ratios of 75/25 and 50/50 were conducive to ryegrass's root growth, which may favor the function of microorganisms for improving the efficiency of wastewater treatment. Like many other plant species (Cramer and Lewis, 1993; Frechilla et al., 2001), ryegrass exhibited

severe reduction in root growth in the 100%  $\text{NH}_4^+$  treatment, implying that ryegrass may not be effective in treating wastewater with sole  $\text{NH}_4^+$  in CWs.

### **Effects of $\text{NO}_3^-/\text{NH}_4^+$ Ratios on Biomass Allocation, Physiological and Morphological Traits**

Not only dry matter production but also the partitioning of dry matter between shoot and root was affected by the inorganic N ratio in the nutrient solution. Ryegrass R/S ratios were lower in  $\text{NH}_4^+$ - than  $\text{NO}_3^-$ -fed plants, similar to findings in other experiments (Cramer and Lewis, 1993; Guo et al., 2002; Tylova et al., 2005). For example, Tylova et al. (2005) reported that R/S of *Phragmites* was on average 27% higher under 100%  $\text{NO}_3^-$  than 100%  $\text{NH}_4^+$  treatment, but was 175% higher for ryegrass in this study, which indicated more depressed root growth of ryegrass under sole  $\text{NH}_4^+$  supply. This again illustrates that ryegrass does not have an advantage in treating wastewater with 100%  $\text{NH}_4^+$ .

Our data also indicate significant effects of N form on leaf traits such as leaf area, chlorophyll concentration and SLA. The response of leaf area to different N forms was consistent with that of plant DM in this study. Sole  $\text{NO}_3^-$  or  $\text{NH}_4^+$  supply had restricted leaf expansion, similar to what Guo et al. (2002) reported for French bean (*Phaseolus vulgaris* L.). Based on treatment effects on leaf area, we can infer treatment effects on plant growth. However, the effects of N form on SLA are complex and depend on plant species, the degree of maturity of the leaf, and the environmental conditions (Bruck and Guo, 2006). In this study, from 100%  $\text{NO}_3^-$  to 100%  $\text{NH}_4^+$  the SLA decreased while the chlorophyll concentration increased, suggesting that ryegrass leaves thickened and concentrated chlorophyll under 100%  $\text{NH}_4^+$  supply. Ruan et al. (2007) reported that tea plants supplied with  $\text{NO}_3^-$  displayed yellowish leaves, consistent with the observed decrease in chlorophyll concentrations in the  $\text{NO}_3^-$  only treatment in this study.

### **Tissue N Concentration and N Accumulation under Different $\text{NO}_3^-/\text{NH}_4^+$ Ratios**

Less energy is required in the process of  $\text{NH}_4^+$  assimilation (as compared with  $\text{NO}_3^-$  assimilation) for higher plants (Zhang et al., 2007). A lower  $\text{NO}_3^-/\text{NH}_4^+$  ratio in the solution led to higher N concentrations in ryegrass, indicating that the increase in  $\text{NH}_4^+$  concentration enhances the N uptake ability per unit biomass although too much  $\text{NH}_4^+$  suppresses plant root growth.

A significantly higher root  $\text{NO}_3^-$  concentration was found when all N was supplied as  $\text{NO}_3^-$ , consistent with those reported by Lewis and Chadwick (1983), and Murphy and Lewis (1987). However,  $\text{NO}_3^-$  concentrations under any of the treatments were lower than the threshold for animal feed

(Niu et al., 2004; Ge et al., 2007), so all harvested biomass can be used as animal feedstock.

Like the performance of wheat (Cox and Reisenauer, 1973), ryegrass fed with  $\text{NO}_3^-/\text{NH}_4^+$  ratios of 75/25 and 50/50 can achieve the most N accumulation due to its high DM yield rather than high N concentration. The tight relationships between plant shoot DM and N accumulation imply that in wastewater treatment N accumulation was manifested by vigorous plant growth but N concentration was little affected and played an insignificant role in N accumulation. Furthermore, it has been experimentally proven that harvesting plant shoots is an effective way to remove N from wastewater treatment systems (Sundaravadivel and Vigneswaran, 2001; Jiang et al., 2004; Ge et al., 2007).

This study provides information for the application of ryegrass in wastewater treatment systems with different  $\text{NO}_3^-/\text{NH}_4^+$  ratios; however, further study considering the role of microorganisms that live in the substrate is needed to better understand the interactions between plants and microorganisms and their joint contribution to nutrient removal from wastewater treatment systems.

## CONCLUSIONS

This hydroponic study showed that ryegrass preferences approximately equal  $\text{NO}_3^-$  and  $\text{NH}_4^+$  supply, with slight preference for a higher  $\text{NO}_3^-$  supply. Ryegrass would be a suitable species to use in CWs to treat wastewater with  $\text{NO}_3^-/\text{NH}_4^+$  ratios between 75/25 and 50/50, as under this condition ryegrass can achieve the highest shoot and root DM, accumulate the highest amount of N, and as a result would be most efficient in removing N from the wastewater by plant harvesting.

## ACKNOWLEDGMENTS

The project was financially supported by the National Natural Science Foundation of China (No. 30870235).

## REFERENCES

- Ali, A., T. C. Tucker, T. L. Thompson, and M. Salim. 2001. Effects of salinity and mixed ammonium and nitrate nutrition on the growth and nitrogen utilization of barley. *Journal of Agronomy and Crop Science* 186: 223–228.
- Britto, D. T., and H. J. Kronzucker. 2002.  $\text{NH}_4^+$  toxicity in higher plants: a critical review. *Plant Physiology* 159: 567–584.
- Bruck, H., and S. W. Guo. 2006. Influence of N form on growth and photosynthesis of *Phaseolus vulgaris* L. *Journal of Plant Nutrition and Soil Science* 169: 849–856.
- Caicedo, J. R., N. P. Van Der steen, O. Arce, and H. J. Gijzen. 2000. Effect of ammonia nitrogen concentration and pH on growth rates of Duckweed (*Spirodela Polyrhiza*). *Water Resources* 34: 3829–3835.

- Chang, J., C. L. Yue, Y. Ge, and Y. M. Zhu. 2004. Treatment of polluted creek water by multifunctional constructed wetland in China's subtropical region. *Fresenius Environmental Bulletin* 13: 545–549.
- Cox, W. J., and H. M. Reisenauer. 1973. Growth and ion uptake by wheat supplied N as  $\text{NO}_3^-$ , or  $\text{NH}_4^+$ , or both. *Plant and Soil* 38: 363–380.
- Cramer, M. D., and O. A. M. Lewis. 1993. The influence of nitrate and ammonium nutrition on the growth of wheat (*Triticum aestivum*) and maize (*Zea mays*) plants. *Annals of Botany* 72: 359–365.
- Frechilla, S., B. Lasa, L. Ibarretxe, C. Lamsfus, and P. M. Aparicio-Tejo. 2001. Pea responses to saline stress is affected by the source of nitrogen nutrition. *Plant Growth Regulation* 35: 171–179.
- Ge, Y., S. P. Li, X. Y. Niu, C. L. Yue, Q. S. Xu, and J. Chang. 2007. Sustainable growth and nutrient uptake of plants in a subtropical constructed wetland in southeast China. *Fresenius Environmental Bulletin* 16: 1023–1029.
- Guo, S., H. Bruck, and B. Sattelmacher. 2002. Effects of supplied nitrogen form on growth and water uptake of French bean (*Phaseolus vulgaris* L.) plants. *Plant and Soil* 239: 267–275.
- Hoagland, D. R., and D. I. Arnon. 1950. The water culture method for growing plants without soil. *California Agricultural Experimental Station Circular* 347: 1–32.
- Hunt, R. 1978. *Plant Growth Analysis*. London: Edward Arnold.
- Jiang, Y. P., Y. Ge, C. L. Yue, and J. Chang. 2004. Nutrient removal role of plants in constructed wetland on sightseeing water. *Acta Ecologica Sinica* 24: 1718–1723.
- Lasa, B., S. Frechilla, C. Lamsfus, and P. M. Aparicio-Tejo. 2001. The sensitivity to ammonium nutrition is related to nitrogen accumulation. *Scientia Horticulturae* 91: 143–152.
- Lewis, O. A. M., and S. Chadwick. 1983. An  $^{15}\text{N}$  investigation into N assimilation in hydroponically grown barley (*Hordeum vulgare* L. cv. Clipper) in response to mixed  $\text{NO}_3^-$  and  $\text{NH}_4^+$  nutrition. *New Phytologist* 95: 635–646.
- Liu, D., Y. Ge, J. Chang, C. H. Peng, B. H. Gu, G. Y. S. Chan, and X. F. Wu. 2009. Constructed wetlands in China: Recent developments and future challenges. *Frontiers in Ecology and the Environment* 7: 261–268.
- Lu, W. X., Y. Ge, J. Z. Wu, and J. Chang. 2004. Study on the method for the determination of nitric nitrogen, ammoniacal nitrogen, and total nitrogen in plant. *Spectroscopy and Spectral Analysis* 24: 204–206.
- Murphy, A. T., and O. A. M. Lewis. 1987. Effect of N feeding sources on the supply of N from root to shoot and the site of N assimilation in maize (*Zea mays* L. cv. R201). *New Phytologist* 107: 327–333.
- Niu, X. Y., Y. Ge, J. Chang, B. H. Guan, and Q. S. Xiu. 2004. The role of *Lolium perenne* in constructed wetland ecological engineering for eutrophic water purification. *Wetland Science* 2: 202–207.
- Peng, Y. S., and E. Liu. 1992. Studies of method on extract chlorophyll a and b. *Acta Agricultural Universitas Pekinensis* 18: 247–250.
- Ruan, J. Y., J. Gerendas, R. Hardter, and B. Sattelmacher. 2007. Effect of nitrogen form and root-zone pH on growth and nitrogen uptake of tea (*Camellia sinensis*) plants. *Annals of Botany* 99: 301–310.
- Stottmeister, U., A. P. Wiefner, P. Kuschik, U. Kappelmeyer, M. Kastner, O. Bederski, R. A. Muller, and H. Moormann. 2003. Effects of plants and microorganisms in constructed wetlands for wastewater treatment. *Biotechnology Advances* 22: 93–117.
- Sundaravadivel, M., and S. Vigneswaran. 2001. Constructed wetlands for wastewater treatment. *Critical Reviews in Environmental Science and Technology* 31: 351–409.
- Tylova, E., B. Lorenzen, H. Brix, and O. Votrubova. 2005. The effects of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  on growth, resource allocation and nitrogen uptake kinetics of *Phragmites australis* and *Glyceria maxima*. *Aquatic Botany* 81: 326–342.
- Zhang, F. C., S. Z. Kang, F. S. Li, and J. H. Zhang. 2007. Growth and major nutrient concentrations in *Brassica campestris* supplied with different  $\text{NH}_4^+/\text{NO}_3^-$  ratios. *Journal of Integrative Plant Biology* 49: 455–462.
- Zhang, Y. P., X. Y. Lin, Y. S. Zhang, S. J. Zheng, and S. T. Du. 2005. Effects of nitrogen levels and Nitrate/ammonium ratios on oxalate concentrations of different forms in edible parts of spinach. *Journal of Plant Nutrition* 28: 2011–2025.